



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON SOME METALLIC DERIVATIVES OF ETHYL
THIOGLYCOLLATE¹

BY CHARLES A. ROUILLER

PHARMACOLOGICAL LABORATORY, JOHNS HOPKINS UNIVERSITY

Communicated by J. J. Abel, March 10, 1919

In 1910 Abel² discovered that thioglycollic ester dissolves antimony trioxide with the greatest ease, forming an antimony derivative, $\text{Sb}(\text{SCH}_2\text{CO}_2\text{C}_2\text{H}_5)_3$, according to the equation: $2\text{Sb}_2\text{O}_3 + 6\text{HSCH}_2\text{CO}_2\text{C}_2\text{H}_5 = 2\text{Sb}(\text{SCH}_2\text{CO}_2\text{C}_2\text{H}_5)_3 + 3\text{H}_2\text{O}$. The antimony compound separates as a heavy oil which, when treated in absolute alcohol with ammonia, yields the corresponding amide, $\text{Sb}(\text{SCH}_2\text{CONH}_2)_3$, obtained by precipitation from alcohol with ether as a colorless or slightly reddish, semiresinous mass soluble in water in all proportions with neutral reaction. The experiments of Rowntree, carried out in collaboration with Abel, showed that the new amide is a very powerful trypanosomicidal substance.

Professor Abel found that the thioglycollic ester reacts energetically with mercuric oxide also, and in order to determine whether the reaction discovered by him is of general applicability, he suggested to the writer that he try the action of various other metallic oxides on the ester. It was hoped that the resulting products might be so slightly soluble as not to be toxic when applied on open wound surfaces, but yet soluble enough to be antiseptic and bactericidal. Abel's expectation that his reaction would prove to be general has been confirmed; whether the products formed are of pharmacological and therapeutic value we have not yet had an opportunity to determine.

Following Abel's general method (for details see the forthcoming paper in the Journal of the American Chemical Society), the compounds listed below have been prepared and analyzed.

Triethyl bismuthtrithioglycollate, $\text{Bi}(\text{SCH}_2\text{CO}_2\text{C}_2\text{H}_5)_3$

Diethyl mercurydithioglycollate, $\text{Hg}(\text{SCH}_2\text{CO}_2\text{C}_2\text{H}_5)_2$

Ethyl silverthioglycollate, $\text{AgSCH}_2\text{CO}_2\text{C}_2\text{H}_5$.

A copper compound with 9.3 % of copper for which no simple formula can be derived; the normal compound, $\text{Cu}(\text{SCH}_2\text{CO}_2\text{C}_2\text{H}_5)_2$, would contain 21.06% of copper.

Zinc, arsenic, and tin compounds were also prepared but have not yet been analyzed.

¹ A more detailed report of this investigation will appear in the May issue of the Journal of the American Chemical Society.

² Rowntree, L. G., and Abel, J. J., *J. Pharmacology Exper. Therapeutics*, **2**, 1910, (108).